

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

94

B

IFW



Attorney Docket
032340 WN 00

P A T E N T

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Confirmation No.: 8475

In re the application of:

Peter Kótay NAGY, et al.

US Serial No. : 09/701,732

Filed : February 9, 2001

Group Art Unit: 1624

Examiner: Bernhardt, E.

For : PROCESS FOR THE PREPARATION OF A 3/2H-PYRIDAZINONE-4-
SUBSTITUTED AMINO-5-CHLORO-DERIVATIVE

RESPONSE TO NOTICE OF DRAWING INCONSISTENCY
WITH SPECIFICATION

Mail Stop Issue Fee

Commissioner for Patents

P.O. Box 1450

Alexandria, VA 22313-1450

Sir:

On June 23, 2004, the U.S. Patent and Trademark Office mailed a NOTICE OF DRAWING INCONSISTENCY WITH SPECIFICATION ("NOTICE"), a copy of which is enclosed, in connection with the above-identified application. According to the NOTICE, Figure 1 is contained in the Drawings but not listed in the Brief Description of the Drawings in the specification. The NOTICE requires correction.

Applicants wish to point out to the Patent Office that the above-identified application does not contain any drawings, i.e., there is no Figure 1. At the direction of a Patent Office representative at the Office of Patent Publication/Publishing Division, Applicants are filing this paper to explain that the specification does not contain any

drawings. Submitted herewith are copies of the specification and the Filing Receipt (which does not indicate the presence of drawings).

Thus, Applicants submit that the above-identified application does not contain a drawing inconsistency with the specification.

Respectfully submitted,

SMITH, GAMBRELL & RUSSELL, LLP

By: Mary A. Montebello (Reg# 33,021)
for Robert G. Weilacher, Reg. No. 20,531.
1850 M Street, N.W., Suite 800
Washington, D.C. 20036
Telephone: (202)263-4300
Facsimile: (202) 263-4329

Dated: July 13, 2004

Enclosures: Copies of

- (1) Notice of Drawing Inconsistency with Specification
- (2) Specification
- (3) Filing Receipt



COPY


UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Drawing Due 7/24/04

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
09/701,732	02/09/2001	Peter Kotay Nagy	32340WC004	8475

7590 06/23/2004
Smith Gambrell & Russell
Beveridge DeGrandi Weilacher & Young
1850 M Street N W Suite 800
Washington, DC 20036



EXAMINER

BERNHARDT, EMILY B

ART UNIT PAPER NUMBER

1624

DATE MAILED: 06/23/2004

Please find below and/or attached an Office communication concerning this application or proceeding.

DOCKET _____
DOCKETED BY: QB
DATE 6/25/04
GIVEN TO RGW/BN



09/701,732

UNITED STATES PATENT AND TRADEMARK OFFICE

UNDER SECRETARY OF COMMERCE FOR INTELLECTUAL PROPERTY AND
DIRECTOR OF THE UNITED STATES PATENT AND TRADEMARK OFFICE

NOTICE OF DRAWING INCONSISTENCY WITH SPECIFICATION

The drawings filed 12/4/00 have been received. However, an inconsistency exists between the drawings and the Brief Description of the Drawings in the specification.

Figures _____ are listed in the Brief Description of the Drawings in the specification but not contained in the Drawings.

Figures 1 are contained in the Drawings but not listed in the Brief Description of the Drawings in the specification.

Applicant is required to correct the above-noted inconsistency within a time period of **ONE MONTH or THIRTY (30) DAYS, whichever is longer**, from the mailing date of this Notice, or within the time remaining in the time period set forth in the Notice of Allowability (Form PTOL-37) to file corrected drawings, whichever is longer. **NO EXTENSION OF THIS TIME PERIOD MAY BE GRANTED UNDER EITHER 37 CFR 1.136 (a) OR (b)**

Failure to correct the above noted inconsistency will result in abandonment of the application.

The file will be held in the Publishing Division to await the correction of the inconsistency.

Return Corrected Drawings/Specification to:

Mail Stop Issue Fee
Commissioner for Patents
P.O. Box 1450


Alexandria, VA 22313-1450

Office of Patent Publication/Publishing Division
Customer Service: 703-308-6789
1-888-786-0101

COPY

PROCESS FOR THE PREPARATION OF A 3(2H)-
PYRIDAZINONE-4-SUBSTITUTED AMINO-5-CHLORO-
DERIVATIVE


The invention relates to a process for the preparation of 5-chloro-4-{3-[N-(2-(3,4-dimethoxyphenyl)-ethyl)-N-methyl-amino]-propylamino}-3(2H)-pyridazinone of the formula (I).



The British patent specification No. 2 262 526 provides new 3(2H)-pyridazinone-4-substituted amino-5-halo derivatives which possess valuable antiarrhythmic properties and prevent ventricular and auricular fibrillations. The 5-chloro-4-{3-[N-(2-(3,4-dimethoxyphenyl)-


ethyl]-N-methyl-amino]-propylamino}-3(2H)-pyridazinone of formula (I) is described in the above-mentioned British patent specification.

According to the British patent specification No. 2 262 526 the compound of formula (I) is prepared by reacting 4,5-di-chloro-3(2H)-pyridazinone of formula (XI)




(XI)

with the amine of formula (X).



(X)

The drawback of the process resides in the fact that a mixture of the desired compound of formula (I) and the regioisomer thereof of formula (IA)




is obtained, wherein the main component is the undesired isomer of formula (IA), while the desired compound of formula (I) is present only as a side-product, in an amount of a few %.

Only by expensive and cumbersome column chromatography can the compound of formula (I) be separated and isolated in a pure state from the thus-obtained mixture. A further disadvantage of the method is that a considerable (2.5-3-fold) molar excess of the expensive amino component of formula (X) obtained in a multi-step reaction is applied, which renders the method less economical.


The present invention aims at providing a more regio-selective method for the preparation of 5-chloro-4-{3-[N-[2-(3,4-dimethoxyphenyl)-ethyl]-N-methylamino]-propyl-amino}-3(2H)-pyridazinone of the formula (I), which is devoid of the drawbacks of the hitherto known processes.

It has been found that the above aim can be achieved by producing the 5-chloro-4-{3-[N-[2-(3,4-dimethoxyphenyl)-ethyl]-N-methylamino]-propylamino}-3(2H)-pyridazinone of formula (I) and pharmaceutically acceptable acid addition salts thereof according to the method of the invention, which comprises

a₁) reacting a compound of the general formula (II),




wherein X stands for a leaving group, with N-methyl-homoveratryl amine of the formula (VI);



or

a₂) reacting a compound of the general formula (III),




wherein R stands for lower alkanoyl, aroyl or aryl-(lower alkanoyl), with an agent containing a leaving group of the formula X and reacting the thus-obtained compound of the general formula (II) with the compound

of formula (VI);

or


- a₃) reacting 4-(3-hydroxypropylamino)-3,5-dichloro-pyridazine of the formula (IV)



with an agent suitable for introducing a group of the formula R, reacting the thus-obtained compound of general formula (III) with an agent containing a leaving group of the formula X and reacting the thus-obtained compound of general formula (II) with the compound of formula (VI);

or


- a₄) reacting 3,4,5-trichloropyridazine of the formula (V)



with 3-amino-1-propanol, reacting the thus-obtained compound of formula (IV) with an agent suitable for introducing a group of the formula R, reacting the thus-obtained compound of general formula (III) with an agent containing a leaving group of the formula X and reacting the thus-obtained compound of general formula (II) with a compound of the formula (VI);


or

- b₁) removing the group of the formula R (wherein R is as stated above) from a compound of general formula (IX);



or


- b₂) reacting the compound of formula (VIII)



with an agent suitable for introducing a group of the formula R and removing the group of formula R from the thus-obtained compound of general formula (IX);

or

b₃) reacting a compound of the general formula (VII),



wherein X is as stated above, with a compound of the formula (VI), reacting the thus-obtained compound of formula (VIII)


with an agent suitable for introducing a group of the formula R, and removing the group of the formula R from the thus-obtained compound of general formula (IX);
or

b₄) reacting the compound of formula (IV) with an agent containing a leaving group of the formula X, reacting the thus-obtained compound of general formula (VII) with the compound of formula (VI), reacting the thus-obtained compound of general formula (VIII) with an agent suitable for introducing a group of the formula R and removing the group of the formula R from the thus-obtained compound of general formula (IX);

and, if desired, converting the thus-obtained compound of formula (I) into an acid addition salt thereof.

The invention is based on the discovery that the regio-selectivity of the reaction can be improved considerably when using 3,4,5-

trichloropyridazine of the formula (V) as starting substance. When reacting the compound of formula (V) with 3-amino-1-propanol an approximately 1:1 mixture of the desired compound of formula (IV) and the regioisomer thereof of formula (IVA)



is obtained. A further advantage of the application of the compound of formula (V) as starting substance resides in the fact that the isomers of formulae (IV) and (IVA) can readily be separated by crystallization, and thus the expensive column chromatography cumbersome on an industrial scale can be eliminated. A further advantage of the process according to the invention is that the regioisomers are separated at the beginning of the synthesis, when the first intermediate is formed, so the further reaction steps and the closing step are

carried out with the application of only one regioisomer. Thus the desired product can be separated from the reaction mixture with a reduced loss and in a higher purity compared to the hitherto known processes. It was not unforeseen that the regioisomers of formulae (IV) and (IVA) can be separated so simply, by crystallization, and converted to the compounds of general formulae (II) and (III) with such a high yield.

In the first step of variant a) according to the invention 3,4,5-trichloropyridazine of the formula (V) is reacted with 3-amino-1-propanol. The reaction is carried out in an organic solvent. As reaction medium preferably lower alkanols (such as methanol, ethanol, n-propanol, preferably ethanol) or dipolar aprotic solvents (such as acetonitrile or dimethylformamide) are used. The reaction is carried out in the presence of an acid binding agent. For this purpose inorganic acid binding agents (e.g. alkali carbonates, such as sodium

carbonate or potassium carbonate, alkali hydrogen carbonates, such as sodium hydrogen carbonate or potassium hydrogen carbonate), or organic acid binding agents (e.g. amines, such as triethylamine or diethyl isopropyl amine) can be used. According to a preferable embodiment of the process according to the present invention the excess of 3-amino-1-propanol used as reactant may serve as solvent. The reaction can be performed at a temperature between 50 C° and 100 C°, preferably at the boiling point of the reaction mixture.

When the reaction has been accomplished the reaction mixture is preferably worked up by removing the solvent and treating the residue with distilled water or with a 5 to 15 % sodium chloride solution. Thus the two isomers can readily be separated, as the precipitate rich in the undesired isomer of the formula (IVA) can be isolated easily, by filtration, from the aqueous solution rich in the desired isomer of

the formula (IV). If desired, both isomers can be subjected to further purification. The isomer of formula (IVA) can be purified by recrystal-lization from an alcohol, while the compound of formula (IV) can be purified by extraction carried out with an organic solvent (e.g. ethyl acetate or halogenated hydrocarbons, such as dichloroethane or chloroform) followed by drying and evaporating the extract and re-crystallizing the residue from diethyl ether.

In the second reaction step of variant a) the thus-obtained compound of formula (IV) is reacted with an agent suitable for introducing a group of the formula R, wherein R is a lower alkanoyl (e.g. acetyl, propionyl or butyryl), aroyl (e.g. benzoyl optionally carrying a substituent selected from the group consisting of halogen, alkoxy and trifluoro-methyl) or aryl-(lower alkanoyl) (e.g. phenylacetyl). Compounds of the general formula (III) containing an acetyl group in the place of R

can be prepared and used advantageously in the synthesis.

The starting compound of formula (IV) applied for the second step of the synthesis may be either purified or unpurified. Surprisingly it has been found that when the compound of the formula (IV) is unpurified, the compound of general formula (III) is obtained in at least such a high purity and good yield than when starting from a purified compound of the formula (IV). If a compound of the general formula (III) containing acetyl in the place of R is prepared, the compound of formula (IV) is reacted with acetic acid, in the presence of an excess of sodium acetate. As reaction medium preferably glacial acetic acid is used and the sodium acetate is applied in a 2.5 to 3-fold molar excess. The reaction may be carried out at a temperature between 80 °C and 120 °C, it is performed preferably at a temperature of about 100 °C. The reaction mixture can be worked up by extraction carried out with an

organic solvent (preferably dichloromethane) followed by drying and evaporating the organic phase. The product is purified by recrystallization from an alkanol (preferably methanol).

The compound of formula (III) obtained in the third reaction step of variant a) is reacted with an agent containing a leaving group of the formula X, wherein X represents preferably a halogen atom (e.g. chlorine or bromine) or an alkylsulfonyloxy (such as benzenesulfonyl-oxy, p-tolylsulfonyloxy or p-bromophenylsulfonyloxy) group.

It is preferable to carry out the reaction via an intermediate of the general formula (II), wherein X stands for bromine. In this case the compound of general formula (III) is reacted with an aqueous hydrogen bromide solution. It is preferable to use a 48 % aqueous hydrogen bromide solution. Thus the group of formula R can be removed from the amino and hydroxy

groups in excellent yields in a single reaction step, and 4-(3-bromopropylamino)-5-chloro-3(2H)-pyridazinone of the formula (II) is obtained. The reaction is carried out at a temperature between 80 °C and 110 °C, preferably at about 98 °C. The reaction mixture can be worked up readily. The separated product is isolated by filtration or centrifugation and optionally crystallized from an alcohol. The compound of general formula (II) containing a bromine atom in the place of X is a highly preferable intermediate, because the bromine atom is a leaving group easy to be split off.

In the next reaction step of variant a) the compound of general formula (II) is reacted with N-methyl-N-[2-(3,4-dimethylphenyl)-ethyl]-amine (N-methyl-homoveratrylamine). The reaction is carried out in a solvent, in the presence of an acid binding agent. As reaction medium preferably dipolar aprotic solvents (such as acetone, acetonitrile or dimethylformamide) may be used. As acid binding agent

inorganic compounds (e.g. alkali carbonates, such as sodium carbonate or potassium carbonate, or alkali hydrogen carbonates, such as sodium hydrogen carbonate or potassium hydrogen carbonate) or organic compounds (e.g. triethyl-amine or dipropylethylamine) may be used. The reaction is carried out at a temperature between 40 C° and the boiling point of the reaction mixture. One can also proceed by applying excess amine of the formula (VI) which may serve as acid binding agent.

The reaction mixture can be worked up by known methods, e.g. it is evaporated and the residue is poured into water, extracted with an organic solvent (such as dichloromethane or ethyl acetate), the organic extract is filtered, dried and purified by crystallization.

In the first step of variant b) of the process according to the invention the compound of formula (IV) is reacted with an agent containing a leaving group of the formula X.

When preparing compounds of the general formula (VII) containing bromine or chlorine in the place of X the compound of formula (IV) is reacted with thionyl bromide or phosphorus oxybromide, or thionyl chloride or phosphorus oxychloride respectively. The reaction is carried out at a temperature between -10°C and 100°C , in an inert organic solvent. As solvent halogenated hydrocarbons (such as dichloromethane, dichloroethane, chloroform, trichloroethylene, chloro-benzene or carbon tetrachloride), dipolar aprotic solvents (such as acetonitrile) or aromatic solvents (such as benzene or toluene) may be used. The compound of general formula (VII), wherein X stands for bromine, can be prepared from the compound of formula (IV) with aqueous hydrogen bromide in organic acids (e.g. acetic or formic acid), at a temperature between 20°C and 150°C as well.

The compounds of general formula (VII), wherein X stands for alkylsulfonyloxy or arylsulfonyloxy, can be prepared by reacting the compound

of formula (IV) with an appropriate sulfonic chloride in an inert solvent, in the presence of an acid binding agent, at a temperature between -20°C and 60°C . As reaction medium halogenated hydrocarbons (such as trichloromethane, dichloroethane, chloroform, trichloroethylene, chlorobenzene or carbon tetrachloride) or aromatic hydrocarbons (e.g. benzene or toluene) may be applied. As acid binding agent organic bases (e.g. triethylamine or pyridine) can be used.

In the next reaction step of variant b) the thus-obtained compound of general formula (VII) is reacted with an amine of the formula (VI). The reaction is preferably carried out in a dipolar aprotic solvent (such as acetone, aceto-nitrile, dimethylformamide) in the presence of an acid binding agent. As acid binding agent inorganic compounds (such as potassium carbonate or potassium hydrogen carbonate) or organic compounds (such as triethylamine) can be used. An excess of the

amine of the formula (VI) may also serve as acid binding agent. The reaction is carried out at a temperature between 10 C° and the boiling point of the reaction mixture. The reaction mixture can be worked up by known methods, e.g. the solvent is removed, the residue is poured onto water, extracted with an organic solvent (such as dichloromethane or diethyl acetate) and the extract is filtered and dried.

The thus-obtained compound of general formula (VIII) is then reacted with an agent suitable for introducing the group of formula R. The reaction is carried out as specified above in connection with the conversion of the compound of formula (IV) into the compound of formula (III). Preferably compounds of the general formula (IX) containing an acetyl group in the place of R are prepared. For this purpose it is preferable to carry out the reaction of the compound of formula (VIII) in glacial acetic acid, in the presence of anhydrous sodium acetate applied in a 1 to 5-fold molar excess,

at a temperature between 40 C° and 140 C°, preferably between 80 C° and 120 C°.

In the last reaction step of variant b) the group of formula R is removed from the compound of general formula (IX). The reaction is preferably carried out with hydrogen bromide, particularly with 48 % aqueous hydrogen bromide.

The thus-obtained compound of general formula (I) is optionally converted into a pharmaceutically acceptable acid addition salt. The salt formation is carried out by methods known per se, with acids generally used in the pharmacological industry. Both inorganic acids (such as hydrogen chloride, hydrogen bromide, phosphoric acid, sulfuric acid etc.) and organic acids (such as maleic, fumaric, citric, malic, lactic, succinic acid etc.) may be applied. It is preferable to prepare the acid addition salt of the compound of formula (I) formed with hydrogen chloride or fumaric acid.

The compound of formula (V) can be prepared by reacting 4,5-dichloro-3(2H)-pyridazinone with phosphorous oxychloride [T. Kuraishi: Pharm. Bull. (Tokyo) 4, 497 (1956)].

The advantages of the process according to the invention compared to the the hitherto known processes are as follows:

- the reaction is significantly more regioselective than the known processes,
- the desired isomer can be separated from the obtained regioisomer by a simple crystallization, thus the complicated column chromatography cumbersome on an industrial scale can be eliminated,
- the regioisomers are separated at an early stage of the synthesis, consequently only one regioisomer is used in the further steps of the process,
- the reaction steps can be carried out in high yields (e.g. the preparation of the compounds of the formulae (II) and (III)),

the end-product of the formula (I) is obtained in a high yield and with high purity.

The invention is further illustrated by the following Examples of non-limiting character:

Example 1

4-(3-hydroxypropylamino)-3,5-dichloropyridazine
(IV) and 5-(3-hydroxypropylamino-3,4-
dichloropyridazine (IVA)

47.93 g (0.261 mole) of 3,4,5-trichloro-pyridazine are dissolved in ethanol and 49.7 ml ($\rho = 0.982 \text{ g/cm}^3$, 0.65 mole) of 3-amino-1-propanol are added to it under stirring. The solution is heated to boiling, boiled for 30 minutes and a sample is taken for TLC (eluent: a 10:10:0.5 mixture of ethyl acetate : acetone : triethylamine, R_f values: (XI)=0.90, (IV)=0.48, (IVA)=0.32, contamination of unknown structure = 0.75). The reaction takes place

generally within 30 minutes and 1 hour, the whole amount of the starting substance is used up. The reaction mixture is then evaporated, 13 g of sodium chloride are dissolved in distilled water and the thus-obtained solution is added to the evaporated mixture under stirring. The reaction mixture is allowed to stand in a refrigerator overnight at 5 C°. The separated crystals are washed with 10 to 12 ml of cold distilled water and the precipitate is dried. Thus 27.7 g (47.7 %) of crude product (IVA) are obtained. M.p.: 150-153 C°. After recrystallization from methanol the melting point rises to 157-158 C°. The physical characteristics will be specified later.

The aqueous mother liquor is extracted 5 times with 200 cm³ each of ethyl acetate, dried over hot magnesium sulfate, filtered on activated carbon and evaporated to dry. The bulk of the residual crude product is the compound of formula (IV).

Yield of the crude product: 28.02 g (48.32 %),

according to HPLC analysis it contains 7 to 8 % of (IVA) and 1 to 2 % of contamination of unknown structure. The crude product is purified by recrystallization from cold diethyl ether in the following way:

300 ml of diethyl ether are added to it in 5 portions and the oily product is stirred at room temperature. The ether solution is decanted on every occasion and fresh ether is used. The ether solutions are combined, evaporated to a volume of 100 ml and the separated crystals are filtered off. Thus 15.6 g (26 %) of compound of the formula (IV) are obtained. M.p.: 65-66 C°. According to HPLC analysis carried out after purification (IVA) < 3.0 %, and (IV) > 97 %. For the elaboration of the HPLC method small amounts of standards have been prepared by column chromatography.

HPLC method:

Column: Ultrasphere SI 3 mm. 75 cm x 4.6 mm.

Eluent: cyclohexane:ethyl acetate (1:1).

Flow rate: 1.0 ml/min.

Detection: UV 254 nm.

Injected volume: 20 ml (0.8 % dilution).

Retention times: 5.13 for compound (IV) and
13.46 minutes for compound (IVA).

The physico-chemical characteristics of 4-(3-hydroxypropyl-amino)-3,5-dichloropyridazine
(IV):

M.p.: 65-66 C^o.

TLC: ethyl acetate:triethylamine = 20:0.5

R_f = 0.36

Analysis for the formula C₇H₉Cl₂NO₃ (222.08):

C	H	Cl	N
Calculated:			
37.86 %	4.09 %	31.93 %	18.92 %

Found:

37.62	4.12 %	31.71 %	18.67 %
-------	--------	---------	---------

IR (KBr) ν cm⁻¹: 3249, 2947, 1591, 1454, 1390,
1353, 1212, 1177, 1124, 1075, 1037, 908, 683,
522, 460.

¹H-NMR (DMSO): δ 8.70 [s, (1H) pyridazine C-6],
6.8 [t, (1H) 4-NH], 4.7 [t, (1H) OH], 3.74 [qa,

(2H) N-CH₂], 3.5 [qa, (2H) CH₂-O-] 1.73 [m,
(2H) C-CH₂-C].

¹³CNMR (DMSO) δ ppm: 150.8, 116.0, 140.1, 114.7
(pyridazine carbon atoms), (60 C-OH), (43.6 NH-
C), (31.9 C-CH₂-C).

Physico-chemical characteristics of the 5-(3-
hydroxypropyl-amino)-3,4-dichloropyridazine
(IVA):

M.p.: 157-158 C°.

TLC: ethyl acetate:triethylamine = 20:0.5

R_f = 0.16

Analysis for the formula C₇H₉Cl₂N₃O (222.08):

C	H	Cl	N
Calculated:			
37.86 %	4.09 %	31.93 %	18.92 %

Found:

37.68	4.11 %	31.77%	18.73 %
-------	--------	--------	---------

IR (KBr) ν cm⁻¹: 3269, 2935, 1568, 1334, 1283,
1224, 1139, 1070, 1043, 861, 830, 795, 661,
540, 514.

^1H -NMR (DMSO): δ ppm: 8.73 [s, (1H) pyridazine C-6]; 7.59 [t, (1H) 5-NH], 4.66 [t, (1H) OH], 3.4-3.6 [m, (4H) $\text{CH}_2\text{-X}$ X=heteroatom], 1.73 [m, (2H) C- CH_2C].

The stereoscopic vicinity of the NH proton at position 5 and the pyridazine proton at position 6 has been proved by a DNOE experiment.

^{13}C NMR (DMSO) δ ppm: 152.1, 143.7, 137.2, 114.4 (pyridazine carbon atoms), (58.4 C-OH), (39.9 C-NH), (31.4 C- $\text{CH}_2\text{-C}$).

Example 2

Preparation of 4-N-acetyl-4-N-(3-acetoxypropyl)-5-chloro-3(2H)-pyridazinone (III)

Method A

A mixture of 3 g (13.5 mmoles) of 4-(3-hydroxypropylamino)-3,5-dichloropyridazine (IV) and 3 g (36.5 mmoles) of anhydrous sodium

acetate is suspended in 30 cm³ of glacial acetic acid, and the mixture is boiled for 3 hours (TLC ethyl acetate:aceton:triethylamine = 10:10:0.5). The starting substance ($R_f=0.48$) is used up. The reaction mixture is then cooled, 100 cm³ of distilled water are added to it and the mixture is extracted 3 times with 50 cm³ each of dichloromethane. The organic phases are combined, dried over magnesium sulfate, filtered with activated coal and evaporated. The crude oily residue is dissolved in 5 cm³ of hot methanol. Upon cooling 4-N-acetyl-4-N-(3-acetoxypentyl)-5-chloro-3(2H)-pyridazinone (III) begins to separate. The separated crystals are filtered and washed successively with cold methanol and ether. Yield: 2.0 g (51.6 %).

Method B

A mixture of 28 g (0.12 mole) of crude compound of the formula (IV) and 28 g (0.34 mole) of anhydrous sodium acetate is suspended in

280 cm³ of glacial acetic acid. The mixture is heated to boiling and the reaction is followed as specified above. The mixture is then cooled, the sodium acetate is filtered off and washed with glacial acetic acid. The mother liquor is evaporated in vacuo. For the complete removal of the acetic acid 2x50 cm³ of toluene are added to the mixture and it is evaporated again. The residue is then dissolved in 100 cm³ of distilled water, the aqueous mother liquor is extracted 3 times with 100 cm³ each of dichloromethane, dried over magnesium sulfate, filtered over activated carbon and evaporated. The residual crude product (29-30 g) is dissolved in 15-20 cm³ of hot methanol, clarified by activated carbon and filtered while hot. The product separates upon cooling. It is filtered and washed successively with cold methanol and cold ether. Yield: 16-20 g (45-50 %).

The physical and chemical characteristics of the compound of formula (III) prepared according to any of methods A and B have been

found identical.

Physical and chemical characteristics of the 4-N-acetyl-4N-(3-acetoxypropyl)-5-chloro-3(2H)-pyridazinone (III):

M.p.: 108-110 °C.

TLC: acetonitrile:methanol = 9:1 R_f = 0.75

Analysis for the formula $C_{11}H_{14}ClN_3O_4$:

C	H	Cl	N
Calculated:			
45.92 %	4.91 %	12.32 %	14.61 %
Found:			
45.63 %	5.01	12.36 %	14.40 %

IR (KBr) ν cm^{-1} : 3400-2800 [pyridazinone ring (NH-CO)], 1729, 1676 (amides), 1593, 1445, 1406, 1362, 1321, 1256, 1205, 1172, 1128, 1099, 1083, 1032, 971, 945, 888, 845, 831, 777, 741, 637, 610, 569, 448, 427.

1H -NMR (400 MHz, $CDCl_3$) δ ppm: 12.8 [s, (1H)]

pyridazinone-NH], 7.97 [s, (1H) pyridazinone C-6H], 4.12 [t, (2H) J=6.5 Hz, -CH₂O], {3.8 [m, (1H) and 3.7 [m, (1H), -N-CH₂}, 2.03 [s, (3H), CH₃], 1.97 [s, (3H) CH₃], 1.89 [m, (2H), C-CH₂-C].

¹³CNMR (400 MHz CDCl₃) δ ppm: 171.07, 169.68, (acetyl-carbonyl carbon atoms), 159.87 (CO-pyridazinone), 138.2, 138.4, 139.9 (carbon atoms of the pyridazinone ring), 62.0 CH₂-O), 44.2 (CH₂-N), 27.4 (CH₂), 21.7 and 20.9 (CH₃ carbon atoms).

The advantage of method B resides in the elimination of the loss of substance during the recrystallization from diethyl ether.

Example 3

4-(3-bromopropylamino)-5-chloro-3(2H)-pyridazinone (II)

30.5 g (0.106 mole) of 4-N-acetyl-4-N-(3-acetoxypropyl)-5-chloro-3(2H)-pyridazinone

(III) are suspended in 136 cm³ of 48 % aqueous hydrogen bromide solution in a flask that may be closed by a Du Pont screw-cap. The reaction mixture is kept at a temperature between 96 C⁰ and 98 C⁰ for 24 hours under stirring [TLC ethyl acetate:acetone:triethylamine = 10:10:0.5 (III) R_f=0.73]. During that time the starting substance is used up. The reaction mixture is then cooled, the separated crystals are filtered and washed with cold-dichloromethane. Yield: 27.3 g (95 %). The crude product is recrystallized from 100 to 110 cm³ of isopropanol. Yield: 20.2 g (73 %).

Physical and chemical characteristics of the 4-(3-bromo-propylamino)-5-chloro-3(2H)-pyridazinone:

M.p.: 116-118 C⁰.

TLC: ethyl acetate-acetone-triethylamine =
= 10:10:0.5 R_f = 0.73

Analysis for the formula C₇H₉BrClN₃O₄ (266.53):

34

C	H	Cl	N
Calculated:			
31.55 %	3.40 %	13.30 %	15.77 %
Found:			
31.74 %	3.45	13.15 %	15.70 %

IR (KBr) ν cm^{-1} : 3183, 2800, 2400, 1545, 1423, 1374, 1324, 1269, 1239, 1214, 1163, 1107, 1037, 936, 819, 750, 572.

^1H -NMR (DMSO) δ ppm: 12.45 [s, (1H) NH-pyridazine], 7.65 [s, (1H) -pyridazine], 6.4 [s, (1H) 4NH], 3.78 [t, (2H), N-CH₂], 3.58 [t, (2H), Br-CH₂], 2.13 [qa, (2H), CH₂].

^{13}C NMR (DMSO) δ ppm: 156.93 (Co-pyridazinone), 139.8, 105.9 (pyridazinone ring carbon atoms), 34.16 (C-NH), 41.88 (C-Br), 31.95 (CH₂).

Example 4

5-chloro-4-(3-[N-[2-(3,4-dimethoxyphenyl)-ethyl]-N-methyl-amino]-propylamino)-3(2H)-pyridazinone (I)

A mixture of 10.66 g (0.04 mole) of 4-(3-bromopropylamino)-5-chloro-3(2H)-pyridazinone (II), 10.0 g (0.05 mole) of N-methyl-homoveratryl amine (VI) and 8 g of potassium hydrogen carbonate is suspended in 80 cm³ of acetone. The reaction mixture is refluxed for 8 to 12 hours. The reaction is followed by TLC (eluent: ethyl acetate: acetone:triethylamine = 10:10:0.5 (II R_f=0.76), (VI R_f=0.14), (I R_f=0.47)). The mixture is filtered while hot, washed with acetone and the mother liquor is evaporated in vacuo. To the residue 50 cm³ of ethyl acetate are added. The possibly separated inorganic substance is filtered off and the filtrate is evaporated again. The residual viscous, oily product (14 to 15 g) is triturated twice with 50 cm³ of hot water in order to remove the unreacted starting

substance of the formula (VI). The warm aqueous solution is decanted. The oily residue is dissolved in methanol and dried over magnesium sulfate. Upon adding a small amount of diisopropyl ether and cooling the mixture white porous substance separates. Thus 9 g (59.0 %) of crude product are obtained. M.p.: 89-90 °C. After recrystallization from diisopropyl ether 7.0 g (46.0 %) of the title compound are obtained.

Physical and chemical characteristics of the 5-chloro-4-{3-[N-[2-(3,4-dimethoxyphenyl)-ethyl]-N-methylamino]-propyl-amino}-3(2H)-pyridazinone:

M.p.: 90-92 °C.

TLC: ethyl acetate-acetone-triethylamine =
= 10:10:0.5

R_f = 0.45

Analysis for the formula C₁₈H₂₅ClN₄O₃ (380.88):

C	H	Cl	N
Calculated:			
56.76 %	6.62 %	9.31 %	14.71 %
Found:			
56.46 %	6.68	9.26 %	14.85 %

IR (KBr) ν cm^{-1} : 3290, 3111, 2940, 2860, 2830, 2780, 2700, 1640, 1610, 1570, 1520, 1445, 1350, 1260, 1240, 1140, 1100, 950, 900, 800, 600.

^1H -NMR (200 MHz, CDCl_3) δ ppm: 1.7 [s, (1H) pyridazinone-NH], 7.52 [s, (1H) pyridazinone-CH], 6.75 [m, (3H) Ar-H], 6.62 [t, (1H), NH], 3.84 and 3.86 [s, (6H), CH_3O], 3.85 [m, (2H), propyl- CH_2], 2.72 and 2.65 [m, (4H) ethyl- CH_2], 2.56 [m, (2H), propyl- CH_2], 2.33 [s, (3H), N- CH_3], 1.80 [m, (2H), propyl- CH_2].

^{13}C NMR (200 MHz CDCl_3) δ ppm: 157.69 (pyridazinone C3), 148.46, 146.94, 132.64, 120.20, 111.73, 110.93 (CH_3O -phenyl-aromatic carbon atoms), 140.41 (pyridazinone C6), 139.85 (pyridazinone C5), 106.8 (pyridazinone C4),

59.43 (C1-propyl), 55.54 and 55.49 (O-CH₃),
54.83 (C2-ethyl), 42.82 (N-CH₃), 41.67 (C3-
propyl), 32.87 (CH₂-Ar), 27.66 (C2-propyl).


Example 5

Preparation of the starting substance

200 g of commercially available 4,5-dichloro-3(2H)-pyridazinone are refluxed in 1500 cm³ of phosphorus oxychloride for 5 hours. Then the excess of phosphorus oxychloride is distilled off in vacuo. The residue is poured onto icy water, the crystalline product is filtered and dried. Yield: 200 g (89 %) of 3,4,5-trichloro-pyridazine. M.p.: 58-60 C⁰.

What we claim is:


1. A process for the preparation of 5-chloro-4-{3-[N-[2-(3,4-dimethoxyphenyl)-ethyl]-N-methylamino]-propylamino}-3(2H)-pyridazinone of the formula (I)




and pharmaceutically acceptable acid addition salts thereof,

which comprises

- a₁) reacting a compound of the general formula (II),




wherein X stands for a leaving group, with N-methyl-homoveratryl amine of the formula (VI);



or


a₂) reacting a compound of the general formula (III),



wherein R stands for lower alkanoyl, aroyl or aryl-(lower alkanoyl), with an agent containing a leaving group of the formula X and reacting the thus-obtained compound of the general formula (II) with the compound of formula (VI);

or

a₃) reacting 4-(3-hydroxypropylamino)-3,5-dichloro-pyridazine of the formula (IV)



with an agent suitable for introducing a group of the formula R, reacting the thus-obtained compound of general formula (III) with an agent containing a leaving group of the formula X and reacting the thus-obtained compound of general formula (II) with the compound of formula (VI);

or


a₄) reacting 3,4,5-trichloropyridazine of the formula (V)



with 3-amino-1-propanol, reacting the thus-obtained compound of formula (IV) with an agent suitable for introducing a group of the formula R, reacting the thus-obtained compound of general formula (III) with an agent containing a leaving group of the formula X and reacting the thus-obtained compound of general formula (II) with a compound of the formula (VI);


or

- b₁) removing the group of the formula R (wherein R is as stated above) from a compound of general formula (IX);



or


- b₂) reacting the compound of formula (VIII)



with an agent suitable for introducing a group of the formula R and removing the group of formula R from the thus-obtained compound of general formula (IX);

or

b₃) reacting a compound of the general formula (VII),



wherein X is as stated above, with a compound of the formula (VI), reacting the thus-obtained compound of formula (VIII) with an agent suitable for introducing a group of the formula R, and removing the

group of the formula R from the thus-obtained compound of general formula (IX);

or

b₄) reacting the compound of formula (IV) with an agent containing a leaving group of the formula X, reacting the thus-obtained compound of general formula (VII) with the compound of formula (VI), reacting the thus-obtained compound of general formula (VIII) with an agent suitable for introducing a group of the formula R and removing the group of the formula R from the thus-obtained compound of general formula (IX);

and, if desired, converting the thus-obtained compound of formula (I) into an acid addition salt thereof.

2. A process according to variant a₃) or a₄) of claim 1, which comprises reacting the compound of formula (IV) with an agent suitable for introducing an acetyl group.

3. A process as claimed in claim 2, which comprises carrying out the reaction with a mixture of acetic acid and sodium acetate.
4. A process as claimed in claim 3, which comprises carrying out the reaction at a temperature between 80 C° and 110 C°.
5. A process according to any of variants a₂) to a₄) of claim 1, which comprises reacting the compound of general formula (III) with an agent containing a halogen atom, an alkylsulfonyloxy or arylsulfonyloxy group, preferably a chlorine or bromine atom or a methanesulfonyloxy, benzyulfonyloxy, p-toluenesulfonyloxy or p-bromophenylsulfonyloxy group.
6. A process as claimed in claim 5, which comprises reacting the compound of general formula (III) with hydrogen bromide.
7. A process as claimed in claim 6, which comprises carrying out the reaction at a

temperature between 80 C° and 110 C°.

8. A process according to variant a₃) or a₄) of claim 1, which comprises reacting the compound of formula (IV) without purification with the agent suitable for introducing a group of the formula R.

9. A process according to any of variants a₁) to a₄) of claim 1, which comprises carrying out the reaction of the compounds of formulae (II) and (VI) in a dipolar aprotic solvent, in the presence of an acid binding agent.

10. A process as claimed in claim 9, which comprises using as solvent acetone, acetonitrile or dimethylformamide and as acid binding agent an alkali carbonate, alkali hydrogen carbonate or an amine, preferably triethylamine or an excess of the reagent of formula (VI).

11. A process according to variant a₄) of claim 1, which comprises carrying out the reaction of the compound of formula (V) with 3-amino-1-propanol in a lower alcohol or a dipolar aprotic solvent, in the presence of an acid binding agent.

12. A process as claimed in claim 11, which comprises using as solvent ethanol, acetonitrile or dimethyl formamide and as acid binding agent an alkali carbonate, alkali hydrogen carbonate, an organic amine, preferably triethylamine or an excess of the 3-amino-1-propanol.

13. A process as claimed in claim 12, which comprises carrying out the reaction at a temperature between 50 C° and 100 C°.

14. The compound of formula (IV).

15. Compounds of the general formula (III), wherein R stands for a lower alkanoyl, aroyl or

aryl-(lower alkanoyl) group.

16. A compound of the general formula (III) according to claim 15, wherein R stands for acetyl.

17. Compounds of the general formula (II), wherein X stands for a leaving group.

18. A compound of the general formula (II) according to claim 17, wherein X stands for bromine.

Data entry by : NGUYEN, SON**Team : OIPE****Date: 03/01/2001**

**LICENSE FOR FOREIGN FILING UNDER
Title 35, United States Code, Section 184
Title 37, Code of Federal Regulations, 5.11 & 5.15**

GRANTED

The applicant has been granted a license under 35 U.S.C. 184, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" followed by a date appears on this form. Such licenses are issued in all applications where the conditions for issuance of a license have been met, regardless of whether or not a license may be required as set forth in 37 CFR 5.15. The scope and limitations of this license are set forth in 37 CFR 5.15(a) unless an earlier license has been issued under 37 CFR 5.15(b). The license is subject to revocation upon written notification. The date indicated is the effective date of the license, unless an earlier license of similar scope has been granted under 37 CFR 5.13 or 5.14.

This license is to be retained by the licensee and may be used at any time on or after the effective date thereof unless it is revoked. This license is automatically transferred to any related applications(s) filed under 36 CFR 1.53(d). This license is not retroactive.

The grant of a license does not in any way lessen the responsibility of a licensee for the security of the subject matter as imposed by any Government contract or the provisions of existing laws relating to espionage and the national security or the export of technical data. Licensees should apprise themselves of current regulations especially with respect to certain countries, of other agencies, particularly the Office of Defense Trade Controls, Department of State (with respect to Arms, Munitions and Implements of War (22 CFR 121-128)); the Office of Export Administration, Department of Commerce (15 CFR 370.10 (j)); the Office of Foreign Assets Control, Department of Treasury (31 CFR Parts 500+) and the Department of Energy.

NOT GRANTED

No license under 35 U.S.C. 184 has been granted at this time, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" DOES NOT appear on this form. Applicant may still petition for a license under 37 CFR 5.12, if a license is desired before the expiration of 6 months from the filing date of the application. If 6 months has lapsed from the filing date of this application and the licensee has not received any indication of a secrecy order under 35 U.S.C. 181, the licensee may foreign file the application pursuant to 37 CFR 5.15 (b).

PLEASE NOTE the following information about the Filing Receipt:

- The articles such as "a," "an" and "the" are not included as the first words in the title of an application. They are considered to be unnecessary to the understanding of the title.
- The words "new," "improved," "improvements in" or "relating to" are not included as first words in the title of an application because a patent application, by nature, is a new idea or improvement.
- The title may be truncated if it consists of more than 600 characters (letters and spaces combined).
- The docket number allows a maximum of 25 characters.
- If your application was submitted under 37 CFR 1.10, your filing date should be the "date in" found on the Express Mail label. If there is a discrepancy, you should submit a request for a corrected Filing Receipt along with a copy of the Express Mail label showing the "date in."
- The title is recorded in sentence case.

Any corrections that may need to be done to your Filing Receipt should be directed to:



COPY

UNITED STATES PATENT AND TRADEMARK OFFICE

COMMISSIONER FOR PATENTS
UNITED STATES PATENT AND TRADEMARK OFFICE
WASHINGTON, D.C. 20231
www.uspto.gov

APPLICATION NUMBER	FILING DATE	GRP ART UNIT	FIL FEE REC'D	ATTY. DOCKET NO	DRAWINGS	TOT CLAIMS	IND CLAIMS
09/701,732	02/09/2001	1624	1070	32340WC004		18	4

FILING RECEIPT

Smith Gambrell & Russell
Beveridge DeGrandi Weilacher & Young
1850 M Street N W Suite 800
Washington, DC 20036

OC000000005807157

Date Mailed: 03/01/2001

Receipt is acknowledged of this nonprovisional Patent Application. It will be considered in its order and you will be notified as to the results of the examination. Be sure to provide the U.S. APPLICATION NUMBER, FILING DATE, NAME OF APPLICANT, and TITLE OF INVENTION when inquiring about this application. Fees transmitted by check or draft are subject to collection. Please verify the accuracy of the data presented on this receipt. If an error is noted on this Filing Receipt, please write to the Office of Initial Patent Examination's Customer Service Center. Please provide a copy of this Filing Receipt with the changes noted thereon. If you received a "Notice to File Missing Parts" for this application, please submit any corrections to this Filing Receipt with your reply to the Notice. When the PTO processes the reply to the Notice, the PTO will generate another Filing Receipt incorporating the requested corrections (if appropriate).

Applicant(s)

Peter Kotay Nagy, Vac, HUNGARY;
Gyula Simig, Budapest, HUNGARY;
Jozef Barkoczy, Budapest, HUNGARY;
Ilona Sztruhar, Budapest, HUNGARY;
Laszlo Balazs, Budapest, HUNGARY;
Imre Doman, Budapest, HUNGARY;
Zoltan Greff, Budapest, HUNGARY;
Zoltan Ratkai, Budapest, HUNGARY;
Peter Seres, Budapest, HUNGARY;
Tamas Karancsi, Budapest, HUNGARY;

Continuing Data as Claimed by Applicant

THIS APPLICATION IS A 371 OF PCT/HU98/00054 06/05/1998

Foreign Applications

If Required, Foreign Filing License Granted 02/27/2001

Title

Process for the preparation of A 3(2H)-pyridazinone-4- substituted amino 5-chloro derivative

Preliminary Class